

MEAN VALUE THEOREM

- (1) Find values of the constants a , b and c for which the graphs of the two functions $f(x) = x^2 + ax + b$ and $g(x) = x^3 - c$, $x \in \mathbb{R}$ intersect at the point $(1, 2)$ and they have the same tangent there.
- (2) Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable. Assume that $f(0) = g(0)$ and $f'(x) \leq g'(x)$, $\forall x \in \mathbb{R}$. Show that $f(x) \leq g(x)$ for $x \geq 0$.
- (3) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable. Assume that $1 \leq f'(x) \leq 2$ for $x \in \mathbb{R}$ and $f(0) = 0$. Prove that $x \leq f(x) \leq 2x$ for $x \geq 0$.
- (4) Use MVT to establish the following inequalities
- (a) $e^x > 1 + x$, $\forall x \in \mathbb{R}$.
 - (b) $\frac{y-x}{y} < \log \frac{y}{x} < \frac{y-x}{x}$ for $0 < x < y$.
 - (c) $\frac{1}{2\sqrt{n+1}} < \sqrt{n+1} - \sqrt{n} < \frac{1}{2\sqrt{n}}$, $\forall n \in \mathbb{N}$.
 - (d) If $e \leq a < b$, then $a^b > b^a$. (Hint: Use part (b)).
 - (e) **Bernoulli's Inequality:** Let $\alpha > 0$ and $h \geq -1$. Then
$$(1+h)^\alpha \leq 1 + \alpha h, \quad \text{for } 0 < \alpha \leq 1,$$
$$(1+h)^\alpha \geq 1 + \alpha h, \quad \text{for } \alpha \geq 1.$$
- (5) Prove that $\frac{\sin x}{x}$ is strictly decreasing on $(0, \pi/2)$.
- (6) Let $f : [0, 1] \rightarrow \mathbb{R}$ be differentiable such that $|f'(x)| < 1$, $\forall x \in [0, 1]$. Show that f has at most one fixed point.
- (7) Let $f : [0, 1] \rightarrow \mathbb{R}$ be differentiable and $f(0) = 0$. Suppose that $|f'(x)| \leq |f(x)|$ $\forall x \in [0, 1]$. Show that $f = 0$.
- (8) Let $f : (0, 1] \rightarrow \mathbb{R}$ be differentiable with $|f'(x)| < 1$. Define $a_n := f(1/n)$. Show that (a_n) converges.
- (9) Let $f : [a, b] \rightarrow \mathbb{R}$ be differentiable and $a \geq 0$. Using Cauchy mean value theorem, show that there exist $c_1, c_2 \in (a, b)$ such that $\frac{f'(c_1)}{a+b} = \frac{f'(c_2)}{2c_2}$.