

Random Vector

Let (\mathcal{S}, Σ, P) be a probability space. A (univariate) random variable describes a numerical quantities of a typical outcome of a random experiment. In many experiments an observation is expressed as a family of several separate numerical quantiles and we may be interested in simultaneously studying all of then together. Consider the following example.

Example 1. *Two distinguishable dice (labelled as D_1 and D_2) are thrown simultaneously. The sample space is $\mathcal{S} = \{(i, j) : i, j \in \{1, 2, \dots, 6\}\}$. For $(i, j) \in \mathcal{S}$ define*

$$X_1((i, j)) = i + j = \text{sum of number of dots on uppermost faces of two dice}$$

and

$$X_2((i, j)) = |i - j| = \text{absolute difference of number of dots on uppermost faces of two dice.}$$

It may be of interest to study numerical characteristics X_1 and X_2 simultaneously. These considerations lead to the study of the function $\underline{X} = (X_1, X_2) : \mathcal{S} \rightarrow \mathbb{R}^2$

Notations.

- We denote by \mathbb{R}^n the n -dimensional Euclidean space, i.e.,

$$\mathbb{R}^n = \{\underline{x} = (x_1, x_2, \dots, x_n) : x_i \in \mathbb{R}, i = 1, 2, \dots, n\}.$$

- For $i = 1, 2, \dots, n$, let $X_i : \mathcal{S} \rightarrow \mathbb{R}$ be any functions. Then the function $\underline{X} = (X_1, X_2, \dots, X_n) : \mathcal{S} \rightarrow \mathbb{R}^n$ is defined as

$$\underline{X}(w) = (X_1(w), X_2(w), \dots, X_n(w)), w \in \mathcal{S}.$$

- For $A \subseteq \mathbb{R}^n$,

$$\underline{X}^{-1}(A) = \{w \in \mathcal{S} : \underline{X}(w) \in A\}.$$

- For $\underline{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, we denote by $(-\infty, \underline{x}]$ the n -dimensional interval

$$(-\infty, \underline{x}] = (-\infty, x_1] \times (-\infty, x_2] \times \dots \times (-\infty, x_n].$$

Definition 2. *A function $\underline{X} : \mathcal{S} \rightarrow \mathbb{R}^n$ is called a random vector (RV) if $\underline{X}^{-1}((-\infty, \underline{x}]) \in \Sigma$, for all $\underline{x} \in \mathbb{R}^n$. That is, $\{w \in \mathcal{S} : X_1(w) \leq x_1, X_2(w) \leq x_2, \dots, X_n(w) \leq x_n\} \in \Sigma$.*

Example 3. *Let $A, B \subseteq \mathcal{S}$. Define $\underline{X} = (X_1, X_2) : \mathcal{S} \rightarrow \mathbb{R}^2$ by*

$$X_1(w) = I_A(w) = \begin{cases} 1, & \text{if } w \in A, \\ 0, & \text{if } w \notin A; \end{cases}$$

and

$$X_2(w) = I_B(w) = \begin{cases} 1, & \text{if } w \in B, \\ 0, & \text{if } w \notin B. \end{cases}$$

Then \underline{X} is an RV if and only if A and B are events. (Prove!)

Theorem 4. *Let $\underline{X} = (X_1, X_2, \dots, X_n) : \mathcal{S} \rightarrow \mathbb{R}^n$ be a given function. Then \underline{X} is a random vector if and only if X_1, X_2, \dots, X_n are random variables.*

Proof. Exercise. □

Remark 5. *If \mathcal{S} is finite or countable and $\Sigma = \mathcal{P}(\mathcal{S})$, then any function $\underline{X} = (X_1, X_2, \dots, X_n) : \mathcal{S} \rightarrow \mathbb{R}^n$ is a random vector.*

Joint Cumulative Distribution Function

Definition 6. Let $\underline{X} = (X_1, X_2, \dots, X_n) : \mathcal{S} \rightarrow \mathbb{R}^n$ be a random vector. The function $F_{\underline{X}} : \mathbb{R}^n \rightarrow \mathbb{R}$, defined by,

$$F_{\underline{X}}(x_1, x_2, \dots, x_n) = P(\{w \in \mathcal{S} : X_1(w) \leq x_1, X_2(w) \leq x_2, \dots, X_n(w) \leq x_n\}), \quad \forall \underline{x} \in \mathbb{R}^n,$$

is called the **joint cumulative distribution function** (joint c.d.f) or the **joint distribution function** (d.f) of the random vector \underline{X} .

The joint distribution function of any subset of random variables X_1, X_2, \dots, X_n is called a marginal distribution function of $F_{\underline{X}}$.

Remark 7. (1) As in the case of random variables, the set $\{w \in \mathcal{S} : X_1(w) \leq x_1, X_2(w) \leq x_2, \dots, X_n(w) \leq x_n\}$ will be denoted by $\{X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n\}$.

(2) In this course, we will mainly study 2- (and sometimes 3-) dimensional random vectors.

(3) Let $\underline{X} = (X, Y) : \mathcal{S} \rightarrow \mathbb{R}^2$ be a random vector. The joint c.d.f. is a map $F_{\underline{X}} : \mathbb{R}^2 \rightarrow \mathbb{R}$, defined by,

$$F_{\underline{X}}(x, y) = P(\{X \leq x, Y \leq y\}).$$

(4) The c.d.f. of X and Y are called a marginal c.d.f. of $F_{\underline{X}}$.

Proposition 8. Let $\underline{X} = (X, Y) : \mathcal{S} \rightarrow \mathbb{R}^2$ be a random vector with joint c.d.f. $F_{\underline{X}}$. Then the marginal c.d.f. of X and Y are given by

$$F_X(x) = \lim_{y \rightarrow \infty} F_{\underline{X}}(x, y) \quad \text{and} \quad F_Y(y) = \lim_{x \rightarrow \infty} F_{\underline{X}}(x, y)$$

Remark 9. Let $(a_1, b_1), (a_2, b_2) \in \mathbb{R}^2$. Then we know that

$$P(a < X \leq b) = P(X \leq b) - P(X \leq a) = F_X(b) - F_X(a).$$

Now,

$$\begin{aligned} & P(a_1 < X \leq b_1, a_2 < Y \leq b_2) \\ &= P(a_1 < X \leq b_1, Y \leq b_2) - P(a_1 < X \leq b_1, Y \leq a_2) \\ &= [P(X \leq b_1, Y \leq b_2) - P(X \leq a_1, Y \leq b_2)] \\ &\quad - [P(X \leq b_1, Y \leq a_2) - P(X \leq a_1, Y \leq a_2)] \\ &= F_{\underline{X}}(b_1, b_2) - F_{\underline{X}}(a_1, b_2) - F_{\underline{X}}(b_1, a_2) + F_{\underline{X}}(a_1, a_2). \end{aligned}$$

Theorem 10. Let $F_{\underline{X}}$ be the joint cumulative distribution function of a random vector $\underline{X} = (X, Y)$. Then

(1) $\lim_{\substack{x \rightarrow \infty \\ y \rightarrow \infty}} F_{\underline{X}}(x, y) = 1.$

(2) $\lim_{y \rightarrow -\infty} F_{\underline{X}}(x, y) = 0$ and $\lim_{x \rightarrow -\infty} F_{\underline{X}}(x, y) = 0.$

(3) $F_{\underline{X}}(x, y)$ is right continuous and nondecreasing in each argument (keeping other argument fixed).

(4) For each $(a_1, b_1] \times (a_2, b_2]$ in \mathbb{R}^2 ,

$$\Delta = F_{\underline{X}}(b_1, b_2) - F_{\underline{X}}(a_1, b_2) - F_{\underline{X}}(b_1, a_2) + F_{\underline{X}}(a_1, a_2) \geq 0.$$

Theorem 11. Let $G : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function which satisfies properties (1) – (4) of Theorem 10. Then there exists a probability space (\mathcal{S}, Σ, P) and a random vector $\underline{X} = (X_1, X_2, \dots, X_n)$ defined on (\mathcal{S}, Σ, P) such that G is the distribution function of \underline{X} .

Example 12. Let $G : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$G(x, y) = \begin{cases} x, & \text{if } 0 \leq x < 1, y \geq 1, \\ y^2, & \text{if } x \geq 1, 0 \leq y < 1, \\ 1, & \text{if } x \geq 1, x \geq 1, \\ 0, & \text{otherwise.} \end{cases}$$

Show that G is not a distribution function of any random vector (X, Y) .

Solution. Clearly G satisfies properties (1) – (3) of Theorem 10.

For $(a_1, b_1] \times (a_2, b_2]$, where $a_1, a_2 \in [0, 1)$, $b_1, b_2 \in [1, \infty)$ and $a_1 + a_2^2 > 1$. Then

$$G(b_1, b_2) - G(a_1, b_2) - G(b_1, a_2) + G(a_1, a_2) = 1 - a_1 - a_2^2 + 0 < 0.$$

Thus, G is not a joint c.d.f. of any random vector.

Example 13. Consider the function $G : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$G(x, y) = \begin{cases} xy^2, & \text{if } 0 \leq x < 1, 0 \leq y < 1, \\ x, & \text{if } 0 \leq x < 1, y \geq 1, \\ y^2, & \text{if } x \geq 1, 0 \leq y < 1, \\ 1, & \text{if } x \geq 1, y \geq 1, \\ 0, & \text{otherwise.} \end{cases}$$

- (1) Show that G is a joint c.d.f. of some random vector (X, Y) .
- (2) Find the marginal c.d.f. of X and Y .

Solution. Clearly $\lim_{\substack{x \rightarrow \infty \\ y \rightarrow \infty}} G(x, y) = 1$. For fixed $x \in \mathbb{R}$, $\lim_{y \rightarrow -\infty} G(x, y) = 0$ and for fixed $y \in \mathbb{R}$, $\lim_{x \rightarrow -\infty} F_X(x, y) = 0$.

We note that if $y < 0$, then $G(x, y) = 0$ for all $x \in \mathbb{R}$. Moreover,

$$G(x, y) = \begin{cases} 0, & \text{if } x < 0, \\ xy^2, & \text{if } 0 \leq x < 1, 0 \leq y < 1, \\ y^2, & \text{if } x \geq 1, \end{cases}$$

and

$$G(x, y) = \begin{cases} 0, & \text{if } x < 0, \\ x, & \text{if } 0 \leq x < 1, y \geq 1, \\ 1, & \text{if } x \geq 1. \end{cases}$$

One can see that for $y \in \mathbb{R}$, $G(x, y)$ is a continuous (and hence right continuous) function of x . Similarly, for each $x \in \mathbb{R}$, $G(x, y)$ is a continuous function of y .

Furthermore, $G(x, y)$ is nondecreasing in each argument keeping other argument fixed.

For $(a_1, b_1] \times (a_2, b_2]$, we need to show that $\Delta = G(b_1, b_2) - G(a_1, b_2) - G(b_1, a_2) + G(a_1, a_2) \geq 0$. We consider the following cases.

- (1) $a_1 < 0$. Then $\Delta = G(b_1, b_2) - G(b_1, a_2) \geq 0$ as G is nondecreasing.
- (2) $a_2 < 0$.
- (3) $0 \leq a_1 < 1, 0 \leq a_2 < 1, 0 \leq b_1 < 1, 0 \leq b_2 < 1$.
- (4) $0 \leq a_1 < 1, 0 \leq a_2 < 1, 0 \leq b_1 < 1, b_2 \geq 1$.
- (5) $0 \leq a_1 < 1, 0 \leq a_2 < 1, b_1 \geq 1, 0 \leq b_2 < 1$.
- (6) $0 \leq a_1 < 1, 0 \leq a_2 < 1, b_1 \geq 1, b_2 \geq 1$.

- (7) $0 \leq a_1 < 1, a_2 \geq 1, 0 \leq b_1 < 1, b_2 \geq 1.$
- (8) $0 \leq a_1 < 1, a_2 \geq 1, b_1 \geq 1, b_2 \geq 1.$
- (9) $a_1 \geq 1, 0 \leq a_2 < 1, b_1 \geq 1, 0 \leq b_2 < 1.$
- (10) $a_1 \geq 1, 0 \leq a_2 < 1, b_1 \geq 1, b_2 \geq 1.$
- (11) $a_1 \geq 1, a_2 \geq 1, b_1 \geq 1, b_2 \geq 1.$

In all these cases verify that $\Delta \geq 0.$

Therefore, $G(x, y)$ is a distribution function of some random vector $(X, Y).$

The marginal c.d.f. of X and Y are respectively

$$F_X(x) = \lim_{y \rightarrow \infty} G(x, y) = \begin{cases} 0, & \text{if } x < 0, \\ x, & \text{if } 0 \leq x < 1, \\ 1, & \text{if } x \geq 1, \end{cases}$$

and

$$F_Y(y) = \lim_{x \rightarrow \infty} G(x, y) = \begin{cases} 0, & \text{if } y < 0, \\ y^2, & \text{if } 0 \leq y < 1, \\ 1, & \text{if } y \geq 1. \end{cases}$$