

# Types of Random Variables: Discrete and Continuous

Let  $(\mathcal{S}, \Sigma, P)$  be a probability space with a random variable  $X : \mathcal{S} \rightarrow \mathbb{R}$ , and let  $(\mathbb{R}, \mathbb{B}_{\mathbb{R}}, P_X)$  be the probability space induced by  $X$ . Let  $F_X$  be the distribution function of  $X$ . It is known that  $F_X$  uniquely determine  $P_X$  and vice-versa. Thus, to study the induced probability space  $(\mathbb{R}, \mathbb{B}_{\mathbb{R}}, P_X)$ , it is sufficient to study the d.f.  $F_X$ .

In this course we will restrict ourselves to two types of random variables: discrete and continuous. In the first case, the RV assumes at most a countable number of values and hence its d.f is a step function. In the later case, the d.f.  $F_X$  is continuous (we will see the definition later).

**Definition 1.** A random variable  $X$  is said to be of discrete type, or simply discrete, if there exists a finite or a countable set  $E_X \subset \mathbb{R}$  such that  $P(\{X = x\}) > 0, \forall x \in E_X$  and  $P(\{X \in E_X\}) = 1$ . The set  $E_X$  is called the support of the RV  $X$ .

**Remark 2.** (1) If  $X$  is any RV with the d.f.  $F_X$ , then  $P(\{X = x\}) = F_X(x) - F_X(x-)$  for every  $x \in \mathbb{R}$ . (Prove!)

(2) From previous lecture, we know that  $F_X$  is discontinuous at  $x \in \mathbb{R}$  if and only if  $F_X(x-) < F_X(x+) = F_X(x)$ . Hence,  $F_X$  has only jump discontinuities and the size of the jump at any point  $x$  of discontinuity is  $P(\{X = x\}) = F_X(x) - F_X(x-)$ .

**Remark 3.** (1) If  $X$  is a discrete type RV with support  $E_X$ , then

$$P(\{X \in E_X\}) = \sum_{x \in E_X} P(\{X = x\}) = \sum_{x \in E_X} (F_X(x) - F_X(x-)) = 1.$$

(2) The d.f.  $F_X$  is continuous at every point of  $E_X^c$ .

**Definition 4.** Let  $X$  be a discrete type random variable with support  $E_X$ . The function  $f_X : \mathbb{R} \rightarrow \mathbb{R}$  defined by,

$$f_X(x) = \begin{cases} P(\{X = x\}), & \text{if } x \in E_X, \\ 0, & \text{otherwise} \end{cases}$$

is called the probability mass function (p.m.f.) of  $X$ .

**Remark 5.** Let  $X$  be a discrete type RV with support  $E_X$ , the d.f.  $F_X$  and the p.m.f.  $f_X$ .

(1)  $f_X(x) > 0, \forall x \in E_X$  and  $f_X(x) = 0, \forall x \notin E_X$ .

(2)  $\sum_{x \in E_X} f_X(x) = 1$ .

(3) For  $A \in \mathbb{B}_{\mathbb{R}}$ , we have

$$\begin{aligned} P_X(A) &= P_X(A \cap E_X) + P_X(A \cap E_X^c) \\ &= P_X(A \cap E_X) \\ &= \sum_{x \in A \cap E_X} f_X(x). \end{aligned}$$

(4) For  $x \in \mathbb{R}$ , we have

$$F_X(x) = \sum_{y \in (-\infty, x] \cap E_X} f_X(y).$$

**Example 6.** Consider the the random variable defined as  $X(w) = c$  for all  $w \in \mathcal{S}$ , where  $c$  is a fixed real number. Then  $P(\{X = c\}) = 1$  and  $E_X = \{c\}$ . Hence,  $X$  is of discrete

type. Its p.m.f. is given by

$$f_X(x) = \begin{cases} 1, & \text{if } x = c, \\ 0, & \text{otherwise.} \end{cases}$$

**Example 7.** Let  $X$  be the indicator function of  $E$ , where  $E$  is an event. Then  $E_X = \{0, 1\}$  and  $P(\{X \in E_X\}) = 1$ . Thus,  $X$  is discrete and its p.m.f. is given by

$$f_X(x) = \begin{cases} P(E^c), & \text{if } x = 0, \\ P(E), & \text{if } x = 1, \\ 0, & \text{otherwise.} \end{cases}$$

**Example 8.** Consider a coin that, in any flip, ends up in head with probability  $\frac{1}{4}$  and in tail with probability  $\frac{3}{4}$ . The coin is tossed repeatedly and independently until a total of two heads have been observed. Let  $X$  denote the number of flips required to achieve this. Then  $P(\{X = x\}) = 0$ , if  $x \notin \{2, 3, 4, \dots\}$ . For  $n \in \{2, 3, 4, \dots\}$ , let  $S_n = \{(w_1, w_2, \dots, w_n) : w_n = H, w_i = H \text{ for one } i \text{ between } 1 \text{ and } n-1, \text{ and } w_j = T, \text{ for } j \neq i\}$ . Now,

$$\begin{aligned} P(\{X = n\}) &= \sum_{(w_1, w_2, \dots, w_n) \in S_n} P(\{(w_1, w_2, \dots, w_n)\}) \\ &= P(\{(w_1, w_2, \dots, w_n)\} | S_n) \\ &= P(\{w_1\})P(\{w_2\}) \cdots P(\{w_n\}) | S_n \quad (\text{since all events are independent}) \\ &= \frac{1}{4} \left(\frac{3}{4}\right)^{n-2} \frac{1}{4} \binom{n-1}{1} \\ &= \frac{n-1}{16} \left(\frac{3}{4}\right)^{n-2}. \end{aligned}$$

Also,  $\sum_{n=2}^{\infty} P(\{X = n\}) = 1$ . Thus,  $X$  is of discrete type with support  $E_X = \{2, 3, 4, \dots\}$  and p.m.f.

$$f_X(x) = \begin{cases} \frac{x-1}{16} \left(\frac{3}{4}\right)^{x-2}, & \text{if } x \in \{2, 3, 4, \dots\}, \\ 0, & \text{otherwise.} \end{cases}$$

The d.f. of  $X$  is

$$\begin{aligned} F_X(x) &= P(\{X \leq x\}) \\ &= \begin{cases} 0, & \text{if } x < 2, \\ \frac{1}{16} \sum_{j=2}^i (j-1) \left(\frac{3}{4}\right)^{j-2}, & \text{if } i \leq x < i+1, i = 2, 3, 4, \dots, \end{cases} \\ &= \begin{cases} 0, & \text{if } x < 2, \\ 1 - \frac{i+3}{4} \left(\frac{3}{4}\right)^{i-2}, & \text{if } i \leq x < i+1, i = 2, 3, 4, \dots \end{cases} \end{aligned}$$

**Definition 9.** A random variable  $X$  with the d.f.  $F_X$  is said to be of continuous type, or simply continuous, if there exists an integrable function  $f_X : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f_X(x) \geq 0$  for every  $x \in \mathbb{R}$  and

$$F_X(x) = \int_{-\infty}^x f_X(t) dt, \quad x \in \mathbb{R}.$$

The function  $f_X$  is called the probability density function (p.d.f.) of random variable  $X$  and the set  $E_X = \{x \in \mathbb{R} : f_X(x) > 0\}$  is called the support of random variable  $X$  (or of p.d.f.  $f_X$ ).

**Remark 10.** Let  $X$  be a continuous type RV with the support  $E_X$ , the d.f.  $F_X$  and a p.d.f.  $f_X$ .

$$(1) \lim_{x \rightarrow \infty} F_X(x) = F_X(\infty) = \int_{-\infty}^{\infty} f_X(t) dt = 1.$$

- (2)  $F_X$  is continuous on  $\mathbb{R}$ . (Prove!) Therefore,  $P(\{X = x\}) = 0 \forall x \in \mathbb{R}$ . In general, for any countable set  $C$ ,  $P(\{X \in C\}) = 0$ .
- (3) Let  $a, b \in \mathbb{R}$  with  $a < b$ . Then

$$P(\{a < X \leq b\}) = F_X(b) - F_X(a) = \int_a^b f_X(t)dt.$$

In general, for any  $B \in \mathbb{B}_{\mathbb{R}}$ , we have  $P(\{X \in B\}) = \int_{-\infty}^{\infty} f_X(t)I_B(t)dt$ , where  $I_B$  is the indicator function of  $B$ .

**Remark 11.** (1) Suppose that the d.f.  $F_X$  of an RV  $X$  is differentiable at every  $x \in \mathbb{R}$ . Then

$$F_X(x) = \int_{-\infty}^x F'_X(t)dt, \quad x \in \mathbb{R}.$$

This implies that  $X$  is of continuous type and we may take its p.d.f to be  $f_X(x) = F'_X(x)$ ,  $x \in \mathbb{R}$ .

- (2) Suppose that the d.f.  $F_X$  of an RV  $X$  is differentiable everywhere except on a countable set  $C$ . Further suppose that

$$\int_{-\infty}^{\infty} F'_X(t)I_{C^c}dt = 1.$$

This again will imply that  $X$  is of continuous type with p.d.f. (Verify!)

$$f_X(x) = \begin{cases} F'_X(x), & \text{if } x \notin C, \\ a_x, & \text{if } x \in C, \end{cases}$$

where  $a_x$ ,  $x \in C$  are arbitrary non negative constants.

- (3) From the previous remark, it is clear that p.d.f. of a continuous random variable need not be unique.
- (4) There are random variables that are neither of discrete type nor of continuous type. Find some examples.

**Example 12.** Let  $X$  be an RV having d.f.

$$F_X(x) = \begin{cases} 0, & \text{if } x < 0, \\ 1 - e^{-x}, & \text{if } x \geq 0. \end{cases}$$

We observe that  $F_X$  is differentiable everywhere except at  $x = 0$ . Let  $C = \{0\}$ . Moreover,

$$\int_{-\infty}^{\infty} F'_X(t)I_{C^c}dt = \int_0^{\infty} e^{-t} = 1.$$

Hence,  $X$  is of continuous type and its p.d.f. is

$$f_X(x) = \begin{cases} 0, & \text{if } x \leq 0, \\ e^{-x}, & \text{if } x > 0. \end{cases}$$